

# Food Safety Scheme Manual

Appendix 5: Use of in-pack pasteurisation to reduce the risk of *Listeria* contamination on ready-to-eat meats

September 2025

FSSM Appendix 5:	: Use of in-pack	pasteurisation
------------------	------------------	----------------

#### Published by the NSW Food Authority

Title: FSSM Appendix 5: Use of in-pack pasteurisation to reduce the risk of *Listeria* contamination in RTE meats First published: August 2019

© State of New South Wales through NSW Food Authority 2025. You may copy, distribute, display, download and otherwise freely deal with this publication for any purpose, provided that you attribute the Food Authority as the owner. However, you must obtain permission if you wish to charge others for access to the publication (other than at cost); include the publication in advertising or a product for sale; modify the publication; or republish the publication on a website. You may freely link to the publication on a departmental website.

Disclaimer: The information contained in this publication is based on knowledge and understanding at the time of writing (September 2025) and may not be accurate, current or complete. The State of New South Wales (including the Food Authority), the author and the publisher take no responsibility, and will accept no liability, for the accuracy, currency, reliability or correctness of any information included in the document (including material provided by third parties). Readers should make their own inquiries and rely on their own advice when making decisions related to material contained in this publication.

3

# Table of Contents

Introduction	4
Possible disadvantages to in-pack pasteurisation	4
Appropriate process time and temperatures are needed	4
Approval to use an in-pack pasteurisation process	7
Example work instruction for in-pack post pasteurisation process	8
References	8
More information	9

#### Introduction

Standard 4.2.3 – Guidelines for the Management of Listeria (Australian Meat Regulators Group, 2019) requires businesses manufacturing packaged ready-to-eat (RTE) meats for sale to implement an effective Listeria sampling plan that covers environmental testing. The objective of the plan is to assess whether the RTE environment is adequately under control with respect to potential contamination of RTE product with Listeria monocytogenes.

However, businesses that apply a post-cooking lethality treatment to the product, such as in-pack pasteurisation, may be exempted from some of the testing requirements. Applying heat, such as immersing packaged cooked meat in hot water can destroy any *Listeria* that may have contaminated the product.

#### Possible disadvantages to in-pack pasteurisation

- Heating might change the meat characteristics (for example colour, texture, presence of excess fluid etc), making it unattractive to consumers,
- The process is not simple for many businesses to do and is time consuming,
- It can be difficult to heat all surfaces of the meat evenly (might get hot spots and cold spots),
- It may not be suitable for sliced packaged product,
- Heating may adversely affect packaging (not all packaging can withstand high temperatures),
  and
- Heating for too long might begin to heat the core of the product (which will make it difficult to cool).

#### Appropriate process time and temperatures are needed

While in-pack pasteurisation can provide a very effective control measure for *Listeria*, the most difficult part is to find the appropriate time and temperature that will kill *Listeria* without changing the look and taste of the packaged product. There are many variables which can affect the rate of heating and how effective the process is (see Table A5.1), which is why the NSW Food Authority advises that in-pack pasteurisation is only suitable for packaged whole primal pieces such as hams and silversides where only the surface has to be heated to destroy *Listeria*.

Table A5.1. Possible sources of variation for in-pack pasteurisation

Characteristics	Comment
Product shape	The size, shape and thickness of the product will affect the evenness of heating, as will overlapping or touching products, which could create cold spots where the heat will take longer to penetrate
Product surface	Meat coated with spices and skin-on poultry could take longer to heat
Product composition	Injected/pumped/pickled products or presence of nitrite
Packaging	The type of packaging (film composition and thickness)
Type of meat	Lean vs fat; beef vs poultry
Hot water tank	Capacity and ability to heat and hold temperature

The goal of pasteurisation is to heat the surface of the meat to at least 72°C for at least one minute. Businesses are encouraged to get expert help and must validate any process they intend to use.

Table A5.2 shows the times and temperatures required to give a 6D reduction of *Listeria* on the surface of RTE meats. The temperature of the water will need to be hotter than this, and there will be variation between different meat products.

The meat's surface temperature can be measured using an infra-red thermometer (for example, Raytek gun) immediately after removing the product from the hot water or by using a data logger or thermometer inserted into the product just below the surface to ensure the time/temperature parameters in Table A5.2 are met.

Once heated, the product must also be chilled quickly (for example by immersion in iced water or blast freezing straight after heating), as this also helps to destroy *Listeria*.

Table A5.2. Process time and temperature to deliver a 6D reduction of *L. monocytogenes* (MLA, 2011)

Meat surface temperature (°C)	Immersion time (minutes)	(seconds)
60	43	1
61	31	39
62	23	17
63	17	8
64	12	36
65	9	16
66	6	49
67	5	1
68	3	41
69	2	43
70	2	0
71	1	28
72	1	5
73	0	48
74	0	35
75	0	26
76	0	19
77	0	14
78	0	10

A 6D process reduces the number of bacteria from 1,000,000 cells to 1 cell.


#### Approval to use an in-pack pasteurisation process

To qualify for an approval to use in-pack pasteurisation, the following must be applied:

- A fixed calibrated temperature probe or data logger must be in place to make sure that the process reaches the correct temperature and maintained for the correct time.
- If the product is pasteurised in the pack, the package cannot be re-opened or re-packaged before it is sold.
- It is suitable for packaged whole primal pieces (for example hams, silversides) or halves where the area of concern is the external meat surface.
- If the product is diced or sliced, the microbiological area of concern is the core of the product. The pasteurisation time and temperature combination must be measured at the core.
- The procedure must be approved by the NSW Food Authority and documented in the food safety program.
- The process must be verified by submitting a packaged product sample for testing for the presence *Listeria* spp every three months.

If you have an approved process for an in-pack pasteurisation, you do not need to conduct environmental swabbing of work surfaces

### Example work instruction for in-pack post pasteurisation process



#### References

Meat & Livestock Australia (2011). Low temperature cooking of meats. Retrieved from <a href="https://www.mla.com.au/contentassets/fec05c41ae7647cda4688e222f65607c/a.mfs.0248\_final\_report.pd">www.mla.com.au/contentassets/fec05c41ae7647cda4688e222f65607c/a.mfs.0248\_final\_report.pd</a> f (PDF, 296 KB)

## More information

- Visit foodauthority.nsw.gov.au
- Email food.contact@dpird.nsw.gov.au
- Phone 1300 552 406

## Copyright and disclaimer statement

© State of New South Wales through NSW Food Authority 2025. The information contained in this publication is based on knowledge and understanding at the time of writing (September 2025). However, because of advances in knowledge, users are reminded of the need to ensure that the information upon which they rely is up to date and to check the currency of the information with the appropriate officer of the Food Authority or the user's independent adviser.